



For more information about the Organization of Teratology Information Specialists or to find a service in your area, call (866) 626-6847 or visit us online at: www.OTISpregnancy.org.

Pesticides and Pregnancy

This sheet talks about the risks that exposure to pesticides can have during pregnancy. With each pregnancy, all women have a 3% to 5% chance of having a baby with a birth defect. This information should not take the place of medical care and advice from your health care provider.

What are pesticides?

A pesticide is a substance used to prevent or destroy insects, fungus, weeds, or other “pests” which can harm crops or people. For example, pesticides are used to reduce the number of mosquitoes, which may carry the West Nile virus and other diseases that can be spread to people. Pesticides sprayed outside with ground equipment or aircraft are referred to as outdoor pesticides. There are many household products that are also considered pesticides, such as cockroach spray, rat poison, or flea collars for pets. These products are known as indoor pesticides.

How can pesticides get into my body?

Pesticides can get into your body by breathing them in, eating them, or getting them on your skin. During the spraying of outdoor pesticides, some of the pesticide may be carried by wind to other areas. Small amounts of outdoor pesticides can also be found in the food and water supply.

What are the most commonly used pesticides?

There are many different kinds of pesticides used in the United States. Some of the most commonly used pesticides are organophosphate pesticides (examples are malathion and naled) and pyrethrin and pyrethroid pesticides (example: Anvil[®]). After Hurricane Katrina, naled and Anvil[®] were

sprayed over much of the Gulf region to control flies and mosquitoes.

Will exposure to pesticides increase the risk to my pregnancy?

Most animal studies with malathion, naled, pyrethrins, and pyrethroids have not shown an increased risk for birth defects. Low birth weight and higher rates of fetal loss were seen at doses that were poisonous to the mother animal.

Two human studies looked at pregnancy outcomes following the repeated spraying of malathion. These studies did not find a consistent increase in birth defects, and there was no effect on birth weight or miscarriage rates. These studies are limited because although malathion was sprayed in the community, the amount of exposure to each individual is not known. There are no human studies with the outdoor use of naled, pyrethrins, and pyrethroids. However, a study of 113 women using a pyrethroid cream to treat head lice did not show an increased risk for birth defects or pregnancy complications.

Some human studies have associated pesticide exposure during pregnancy with an increased risk of childhood cancer. These studies are limited because the actual amount of pesticide each individual was exposed to is unknown. Other studies have not shown that childhood cancer is related to pesticide exposure. At this point, there is not enough information to determine if exposure to

pesticides during pregnancy increases the risk of childhood cancer.

How can I minimize my exposure to pesticides?

The amount of pesticide is important in deciding whether a pregnancy is at an increased risk. In animal studies, pesticides have generally not been associated with birth defects or pregnancy complications unless the amount of pesticide was high enough to be poisonous to the mother animal. This suggests that direct exposure to pesticides (from mixing and applying them yourself) may pose more risk to a pregnancy than low-level or indirect exposures. As with all chemicals, exposure can be minimized by working in a well ventilated area and using protective equipment. It is unlikely that having your home or workplace treated by a professional exterminator will result in a high enough exposure to increase the risk to a pregnancy.

What if I am exposed to pesticides while breastfeeding?

Pesticides can be found in breast milk, but studies on the effects of these pesticides have not been done. When the milk from nine breastfeeding women living in an area repeatedly sprayed with malathion was tested, no malathion was detected in any of the milk samples. The nutritional and psychological benefits of breastfeeding may outweigh the risk of exposure from pesticides sprayed in your home, workplace, or community.

What if my partner is exposed to pesticides at home or at work?

There are adult safety levels for pesticide exposure that should be followed in the workplace. In animal studies, high doses of malathion reduced male fertility. On the other hand, one human study compared men with high and low sperm counts and did not find a

difference in the amounts of a malathion breakdown product in their bodies. In general, environmental exposures in men have not been associated with birth defects in offspring, although further studies are needed.

December 2005.

Copyright by OTIS.

Reproduced by permission.



The Development of this fact sheet was supported by funds from the Centers for Disease Control and Prevention (CDC).

References:

- Buckley JD, et al. 2000. Pesticide exposures in children with non-Hodgkin lymphoma. *Cancer* 89(11):2315-21.
- Contreras HR, et al. 1999. Morphological alterations in mouse testis by a single dose of malathion. *J Exp Zool* 284:355-9.
- Environmental Protection Agency. Spray drift of pesticides. Retrieved December 2005, from: <http://www.epa.gov/pesticides/factsheets/spraydrift.htm>
- Environmental Protection Agency. What is a pesticide? Retrieved December 2005, from: <http://www.epa.gov/pesticides/about/index.htm>
- Eskenazi B, et al. 2004. Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 112:116-1124.
- Grether JK, et al. 1987. Exposure to aerial malathion application and the occurrence of congenital anomalies and low birthweight. *Am J Public Health* 77:1009-1010.
- Gupta P. 1990. Teratogenic effects of cypermethrin in rats. *J Environ Biol* 11(2):121-126.
- Kennedy D, et al. 2005. Pregnancy outcome following exposure to permethrin and use of teratogen information. *Am J Perinatol* 22(2):87-90.
- Khera KS, et al. 1979. Teratologic assessment of maleic hydrazide and daminozide, and formulations of ethoxyquin, thiabendazole and naled in rats. *J Environ Sci Health B* 14(6):53-77.
- Khera K, et al. 1982. Teratogenicity study on pyrethrum and rotenone (natural origin) in pregnant rats. *J Toxicol Environ Health* 10:111-119.
- Ma X, et al. 2002. Critical windows of exposure to household pesticides and risk of childhood leukemia. *Environ Health Perspect* 110(9):955-60.
- Machin MG and McBride WG. 1989. Teratological study of malathion in the rabbit. *J Toxicol Environ Health* 26(3):249-53.
- Meinert R, et al. 1996. Childhood leukaemia and exposure to pesticides: results of a case-control study in northern Germany. *Eur J Cancer* 32A(11):1943-8.
- Miyamoto J. 1976. Degradation, metabolism and toxicity of synthetic pyrethroids. *Environ Health Criter* 14:15-28.
- Moses M. 1993. Pesticides: in Occupational and Environmental Reproductive Hazards: A Guide for Clinicians. Paul, M (ed). Williams & Wilkins: 296-305.
- Pflieger-Bruss, et al. 2004. The male reproductive system and its susceptibility to endocrine disrupting chemicals. *Andrologia* 36:337-345.
- Reynolds P, et al. 2005. Agricultural pesticide use and childhood cancer in California. *Epidemiology* 16(1):93-100.
- Rueber MD. 1985. Carcinogenicity and toxicity of malathion and malaoxon. *Environ Res* 37(1):119-53.
- Thomas DC, et al. 1992. Reproductive outcomes in relation to malathion spraying in the San Francisco Bay Area, 1981-1982. *Epidemiology* 3(1):32-39.
- Sonawane B. 1995. Chemical contaminants in human milk: an overview. *Environ Health Perspect* 103(Suppl 6):197-205.
- Swan SH, et al. 2003. Semen quality in relation to biomarkers of pesticide exposure. *Environ Health Perspect* 111:1478- 1483.

*If you have questions about the information on this fact sheet or other exposures during pregnancy, call **OTIS** at 1-866-626-6847.*